



Stick or grip? Co-evolution of adhesive toepads and claws in Anolis lizards

Citation

Crandell, Kristen E., Anthony Herrel, Mahmood Sasa, Jonathan B. Losos, and Kellar Autumn. 2014. "Stick or Grip? Co-Evolution of Adhesive Toepads and Claws in Anolis Lizards." *Zoology* 117 (6) (December): 363–369. doi:10.1016/j.zool.2014.05.001.

Published Version

doi:10.1016/j.zool.2014.05.001

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:20444690>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Open Access Policy Articles, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#OAP>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

Stick or grip? Co-evolution of adhesive toepads and claws in Anolis lizards

Kristen E. Crandell¹, Anthony Herrel^{2,3}, Mahmood Sasa^{4,5}, Jonathan B. Losos⁶, Kellar Autumn⁷

4

1. Field Research Station at Fort Missoula, Division of Biological Sciences, University of Montana, Missoula, MT 59812, USA

email : kristen.crandell@umontana.edu

2. UMR 7179 C.N.R.S./M.N.H.N., Département d'Ecologie et de Gestion de la Biodiversité, 57 rue Cuvier, Case postale 55, 75231, Paris Cedex 5, France.

3. Ghent University, Evolutionary Morphology of Vertebrates, K.L. Ledeganckstraat 35, B-9000 Gent, Belgium

4. Palo Verde Biological Station, Organization for Tropical Studies, 0676-2050 San Jose, Costa Rica

5. Instituto Clodomiro Picado, Universidad de Costa Rica, San Jose, Costa Rica

6. Museum of Comparative Zoology and Department of Organismic and Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, MA 02138, USA

7. Department of Biology, Lewis & Clark College, Portland, OR 97219, USA

Pages: 16

Figures: 6

Tables: 1

Key Words: *Anolis*; adhesion; claw; toepad; ecomorphology; clinging; mainland

22

23 ABSTRACT

24 Exploring the relationship between phenotype and performance in an
25 ecological and evolutionary context is crucial to understand the adaptive nature of
26 phenotypic traits. Despite their ubiquity in vertebrates, few studies have examined
27 the functional and ecological significance of claw morphologies. Here we examine
28 the adhesive toepad and claw system of *Anolis* lizards. Claw characters are
29 significantly different between lizards classified as arboreal (perch height > 1 m)
30 and non-arboreal (perch height < 1 m). Arboreal species possess significantly higher
31 and longer claws, and show trends toward decreased claw curvature and wider claw
32 tip angles. Toepad size and claw length and height are tightly correlated with each
33 other and with perch height, suggesting that the adhesive toepad and gripping claw
34 have co-evolved to accommodate different habitats. The functional morphology and
35 evolution of claws are ripe areas for future investigation.

36 INTRODUCTION

37 Studying the link between phenotype and performance is necessary for
38 insight in to the adaptation of form and function (Arnold, 1983). The functional
39 significance of claws is not well studied, despite their ubiquity throughout
40 vertebrates. Claws are known to increase available habitat (Cartmill, 1974), yet the
41 link between morphological and habitat use variation is unknown. Here, we explore
42 the claw in relation to a second structure used in attachment, the adhesive toepad,
43 in *Anolis* lizards.

44 Anoles are an ideal model organism with which to unravel potential
45 ecological contributions of claws, as they occupy a vast array of habitats with
46 predictable evolutionary trajectories (Losos, 2009). *Anolis* lizards comprise a
47 diverse clade that has undergone adaptive radiation on the Caribbean islands of the
48 Greater Antilles, producing species of similar ecology and morphology termed
49 “ecomorphs” that have evolved independently on each island (Williams, 1983).
50 These lizards have been the focus of many studies examining the relationships

1
2
3
4 51 between morphology, performance, and ecology (see Losos, 2009 for a review). A
5
6 52 particularly striking feature of anoles is the adhesive toepad, which enables the
7
8 53 animal to move across smooth substrates with little difficulty. Toepads are thought
9
10 54 to be a key innovation in anoles because their evolution may have allowed these
11
12 55 lizards to occupy a larger portion of the available habitat than previously possible,
13
14 56 thus permitting their radiation and diversification (Warheit et al., 1999).

15
16
17 57 Toepads are effective in permitting anoles to expand their habitat use due to
18
19 58 their attachment abilities: they function best on smooth surfaces, such as leaves and
20
21 59 smooth tree bark. Microscopic hair-like structures on the ventral pad, termed setae,
22
23 60 adhere to substrates via van der Waals forces (Autumn et al., 2002; Puthoff et al.,
24
25 61 2010). Among anole species, clinging ability varies with habitat use. Species that
26
27 62 occur higher in the tree canopy possess larger toepads, and are capable of producing
28
29 63 greater shear forces (Elstrott and Irschick, 2004; Macrini et al., 2003). This variation
30
31 64 suggests that the evolution of adhesive toepads may have been critical for
32
33 65 occupation of arboreal habitats, and thus may have played a major role in the
34
35 66 diversification of Caribbean anoles into a distinct set of ecomorphs.

36
37 67 An often-neglected feature of *Anolis* relevant to clinging is their claws. Like
38
39 68 almost all other lizard species, anoles possess claws, and variation in claw
40
41 69 morphology may be related to differences in habitat use. Unfortunately, claws have
42
43 70 been overlooked not only in anoles, but also in most amniotes. Vertebrates with
44
45 71 claws can occupy larger portions of the habitat than non-clawed animals (Cartmill,
46
47 72 1974). However, the functionality of claws is less understood (see Maddin and Reisz,
48
49 73 2007 and Zani, 2000). Some aspects of claw shape are known to affect attachment
50
51 74 ability, including claw height (the distance measured from dorsal to ventral at the
52
53 75 base of the claw Zani, 2000). In animals such as beetles, claws interact with surface
54
55 76 irregularities in two ways: interlocking and friction. When surface irregularities are
56
57 77 larger than claw tip diameter, the claw mechanically interlocks with the
58
59 78 irregularities. In contrast, when surface irregularities are smaller than claw tip
60
61 79 diameter, attachment results from frictional forces. In this instance, if the tangent

1
2
3
4 80 between the claw and the irregularity is too low, the claw slips (Dai et al., 2002). As
5
6 81 such, mechanical interlocking (accomplished with a smaller claw tip relative to
7
8 82 substrate roughness) is often times stronger than frictional attachment, with a
9
10 83 lower likelihood of failure. Thus, the ability to create a mechanical attachment,
11
12 84 rather than a friction-based attachment, is improved with smaller claw tips.
13
14 85 Decreasing the size (or effective angle) of the tip increases the likelihood of surface
15
16 86 irregularities being larger, offering more opportunity for mechanical attachment. In
17
18 87 fact, in artificial claws, decreasing the tip's effective angle significantly improves the
19
20 88 strength of attachment (Provancher et al., 2004).

21
22 89 However, understanding the broader relationship between claw morphology
23
24 90 and habitat use is not yet possible. A study linking claw morphology directly to
25
26 91 habitat use in birds indicates that ground-dwelling species have significantly less
27
28 92 curved claws than perching species, and species that climb have claws with higher
29
30 93 curvature than both ground and perch dwellers (Feduccia, 1993). Similar patterns
31
32 94 have been observed in lizards: arboreal and saxicolous species have claws with
33
34 95 higher curvature (Tulli et al., 2009). Some additional evidence for the importance of
35
36 96 claws exists in a single species, the ecologically variable *Anolis cybotes*; individuals
37
38 97 in more rocky areas appear to have more curved claws (Wollenberg et al., 2013).

39
40 98 It is reasonable to predict that the claw and adhesive toepad operate under
41
42 99 disparate conditions. The toepad functions best on relatively smooth, homogenous
43
44 100 surfaces. The attachment between microscopic hair-like structures of the ventral
45
46 101 pad and the surface is modeled to be maximal on smoother surfaces (Persson and
47
48 102 Gorb, 2003; Russell and Johnson, 2013), and performance declines with increasing
49
50 103 surface roughness (Vanhooydonck et al., 2005). In contrast, claws appear to
51
52 104 maximize functionality on very rough surfaces, where mechanical interlocking of
53
54 105 the claw is often-times stronger than friction forces on smoother surfaces (Dai et al.,
55
56 106 2002). Thus, it appears both structures are optimal in two different scenarios. This
57
58 107 observation was first put forth by Mahendra (1941), who observed following claw
59
60 108 removal, *Hemidactylus* geckos were unable to attach to rough surfaces, but
61
62
63
64
65

attachment to smooth surfaces was unaffected. Further recent explorations are limited, but best evidenced in an analysis of performance and morphology. Zani (2000) found that in 85 species of lizard, toe width and lamellae number is correlated with attachment ability to smooth surfaces, while claw height is correlated with rough surface attachment.

Examining both the adhesive toepad alongside the claw may reveal ecological patterns hitherto unknown, or previously overlooked and attributed to toepad function alone. Here, we explore the relationship between toepad clinging ability, claw morphology, and habitat use in *Anolis*. We test for associations between claw morphology and habitat use. We predict that arboreal species with higher perch heights will have claw morphologies associated with improved attachment abilities: higher and longer (Zani, 2000), more curved (Feduccia, 1993), and sharper tips (Dai et al., 2002). We also test if phylogenetically corrected features of claws that are thought to improve attachment co-vary with toepad adhesion ability, given that toepads also correlate with habitat (Elstrott and Irschick, 2004). Our available lizards are mainland species, and as such we also confirm that mainland species follow the same trends as island species by testing for a positive relationship between toe force production and perch height (Irschick et al., 1997; Macrini et al., 2003).

METHODS

Study area and species

Fieldwork occurred at La Selva Biological Station, Playa Piro, and Palo Verde Biological Station in Costa Rica, and Gamboa, Panama. A total of 12 species was studied (see table 1). To capture animals and determine perch height, we walked along trails daily when lizards were active and recorded data for any individual sighted. Perch height was measured as distance from the ground to the individual's

original position, and perch diameter was measured at the location the lizard was first observed. Lizards were captured by hand or noose and kept for no longer than 48 hours. Lizards were kept in one-gallon plastic bags, transported to the field lab and returned to their original location following performance trials.

Adhesion performance

Following capture, one person (KEC) took shear-force measurements. A dual-range force sensor (Vernier) was attached to a vertical acetate covered glass microscope slide with a custom plexiglass attachment and butterfly clip. For each subject, the fourth (longest) digit was isolated and gently applied to the acetate sheet (Fig. 1). These performance measures only reflect attachment ability of the adhesive toepad, not the claw, because the claw was not able to penetrate the acetate coating. The anole was pulled by hand at an approximately constant speed, and the shear force was recorded at 40 Hz. Each subject underwent three repeated trials for left and right fourth toe.

Morphology

For all species in this study, we measured toepad and claw morphological characters from preserved specimens at the Museum of Comparative Zoology, Harvard University, and specimens from the Museum of Southwestern Biology, University of New Mexico (see appendix for list). A scanner (Epson Perfection 4900 and V500) digitized images of the fourth right hind digit, with the claw flattened sagittally against the scanner. Measurements were taken with ImageJ (1.4g, Rasband). Morphological characters of the toe included toepad area, measured from where the pad begins to widen (i.e., where the next most distal lamellae is longer than the previous), and lamellae number, counted as lamellae contained within toepad area. Measurements of the claw included: height, length, curvature (as measured by Zani, 2000), and tip angle (Fig. 2).

163 *Analysis*

164 For all species, we determined descriptive statistics for all continuous
165 characters. To meet assumptions of normality and homoscedasticity, all
166 measurements except for toepad force and claw tip angle were log-10 transformed
167 prior to analysis. We corrected for size in characters that significantly correlated
168 with SVL. To do so, we regressed log-10 adjusted values against log-10 adjusted SVL
169 and calculated residuals, which were used for subsequent regressions.

170 A bivariate Pearson correlation on the log-10 transformed data was used to
171 determine correlations among traits. Log-10 transformed data were then used as
172 input for an independent contrasts analysis (Felsenstein, 1985) using the GEIGER
173 package in R (Harmon et al., 2008). The most complete phylogenetic tree of *Anolis*
174 (Nicholson et al., 2005) served as the basis for analyses (Fig. 3). Panamanian species
175 described in Castañeda and de Queiroz (2013) were added as sister taxa, and *Anolis*
176 *apletophallus* was placed as sister taxon to *Anolis limifrons* (S. Poe, pers. comm.).
177 Branch lengths of the phylogeny were unknown, and were set to arbitrary lengths
178 using a Grafen transformation (Grafen, 1989). The calculated residual values of
179 morphological, performance, and habitat use variables were input into a linear
180 regression analysis to determine correlation coefficients. An analysis of variance
181 (ANOVA) tested for statistical significance for all regressions. All analyses presented
182 account for phylogeny.

183 We also compared claw variables between two groups of anoles classified as
184 “arboreal” or “non-arboreal.” Arboreal species were defined as those with an
185 average perch height greater than one meter, whereas non-arboreal were those
186 found perching below one meter. Species with mean perch heights less than one
187 meter are generally seen on or near the ground, whereas those above one meter are
188 those that are often seen in the canopy. Claw morphologies (claw curvature, claw tip
189 angle, claw height, and claw length) were first compared with a phylogenetically
190 corrected MANOVA. Each character was then compared between groups with a
191 phylogenetically corrected ANOVA. One-tailed tests were used in all comparisons, as

we had a-priori predictions for claw curvature (Feduccia, 1993), claw tip angle (Dai et al., 2002), and claw height and length (Zani, 2000). It should be noted that Zani (2000) found the correlation between claw length and clinging ability on rough surfaces was not robust to changes in phylogenetic branch lengths in one out of four models. Both phylogenetic analyses used the GEIGER package in R (Harmon et al., 2008).

RESULTS

Morphology and performance

Table 1 lists the mean \pm SE for all morphological variables and toepad force production. All variables were size corrected, with analyses performed on the residuals of the data regressed against body size (see methods), with the exception of claw tip angle ($p=0.68$, $r^2=0.02$) and claw curvature ($p=0.91$, $r^2=0.001$), because they were uncorrelated to overall body size. All correlations are the result of phylogenetically independent contrasts to account for ancestral relatedness.

Adjusted toepad area correlates positively with the size adjusted variables claw height ($p<0.001$, $r^2=0.65$, d.f.=1,10), claw length ($p=0.03$, $r^2=0.35$, d.f.=1,10), and non-size adjusted claw curvature ($p=0.05$, $r^2=0.26$, d.f.=1,10). Pad area also correlates positively with toepad force production following corrections for size ($p<0.001$, $r^2=0.75$, d.f.=1,10). Toepad lamella number exhibits similar relationships, correlating positively with claw height ($p=0.02$, $r^2=0.42$, d.f.=1,10), claw length ($p=0.03$, $r^2=0.38$, d.f.=1,10), and toepad force production ($p<0.001$, $r^2=0.83$, d.f.=1,10).

A positive correlation also occurs between toepad force production and claw height ($p=0.01$, $r^2=0.51$, d.f.=1,10) and claw length ($p=0.02$, $r^2=0.46$, d.f.=1,10) (Fig. 4). Force production and claw curvature are also related, but the relationship is not statistically significant ($p=0.08$, $r^2=0.30$, d.f.=1,9). No relationship is present between force production and claw tip angle ($p=0.98$, $r^2<0.0001$, d.f.=1,10).

Habitat use

A positive correlation exists between perch height and toepad shear-force ($p=0.006$, $r^2=0.58$, $d.f.=1,9$) (Fig. 5), but no correlation occurs between adjusted force production and perch diameter ($p=0.22$, $r^2=0.16$, $d.f.=1,9$). Perch height does not linearly correlate with claw variables, including claw height ($p=0.43$, $r^2=0.07$, $d.f.=1,9$), claw length ($p=0.26$, $r^2=0.14$, $d.f.=1,9$), claw curvature ($p=0.56$, $r^2=0.04$, $d.f.=1,9$), and claw tip angle ($p=0.90$, $r^2=0.002$, $d.f.=1,9$).

We also compared claw variables with binomially categorized habitat by dividing anoles in to two classes: “arboreal,” with average perch heights of over one meter, and “non-arboreal,” with average perch heights below one meter. We found general trends in all claw morphologies between our arboreal categories (phylogenetic MANOVA $p=0.054$). Claw tip angle was close to statistically different between categories ($p=0.054$, Fig. 6A), as was claw curvature ($p=0.054$, Fig. 6B). Both claw height ($p=0.007$, Fig. 6C) and claw length ($p=0.029$, Fig. 6D) were statistically significantly different between groups.

234

DISCUSSION

Our study is one of the first to examine toe and claw morphology in a vertebrate with an eye towards performance in a natural environment. We find support for both of our hypotheses. First, the adhesive toepad of *Anolis* species co-varies with claw morphology. Secondly, all claw characters measured are associated with habitat use.

We found strong evidence for co-evolution between adhesive toepads and claws in size-corrected correlations between toepad size and claw height and length (Fig. 4). Adhesive toepads and claws provide clinging capability in different substrate conditions: smooth and rough, respectively. Arboreal animals encounter both types of surfaces as they move on leaves, a smooth substrate, and woody

vegetation, a rough substrate; hence, it is not surprising that more arboreal species, needing greater attachment ability, should have better developed claws and toepads.

We found strong trends and significant differences between species grouped as arboreal (mean perch height > 1 m) and non-arboreal (mean perch height < 1 m). A phylogenetic MANOVA of all claw morphologies came very close to statistical significance at $p = 0.054$, indicating that claw characters are likely related to perch height. With phylogenetic ANOVAs for each character, we were able to dissect this trend. Claw curvature and claw tip angle were not statistically significant between groups, but both showed major trends in that direction, with p values very close to, but slightly above, 0.05. This trend does not fall in line with increased claw curvature in arboreal birds in comparison to ground-dwelling birds (Feduccia, 1993). Given that claw sharpness (measured here as a smaller tip angle) improves attachment ability (Dai et al., 2002; Provancher et al., 2004), it is also surprising that arboreal species trend toward less pointed claw tips. We predict with larger sample sizes, a significant trend would appear. These differences suggest further studies examining claw function are vital. Both claw height and length were significantly different between the two groups (Fig. 5). Claw height improves attachment ability on rough surfaces (Zani, 2000). Interestingly, adhesive pad area is also correlated with perch height (Elstrott and Irschick, 2004). Together, differences in claw shape and toepad function in comparison to perch height serve as further evidence for the co-evolution between claws and pads.

Despite strong differences between arboreal and non-arboreal morphologies, we did not find linear correlations between claw characters and perch height in this study. Arboreality implies a heightened need for both attachment systems, to maintain attachment during motion and to avoid falling. However, the functionality of these systems differs in their basic attachment mechanics. In particular, the toepad area is continuous with an increase in surface area of the pad resulting in an increase in clinging ability. In contrast, claw function may not be continuous – once a certain threshold is reached in size or shape, an increase in size or further changes

in shape may confer no more functional, and therefore selective, advantage. For example, higher claws may serve no better than slightly lower ones, as both are capable of mechanical interlocking with the substrate. This is in sharp contrast with toepad area, where increased size confers a linear increase in force. Thus, certain morphologies will be able to accommodate rough substrates, without regard to the actual perch height.

Our study focused on the two ecological standards for anoles: perch height and diameter. While we found several intriguing patterns with perch height, none of our measured variables correlated with perch diameter. In a study by Macrini et al. (2003), perch diameter in mainland and island anoles correlates with pad area. However, we found no such relationship here. Mainland species vary in habitat use, and with our relatively small sample size, such patterns may not appear due to higher ecological variation of mainland anoles (Schaad and Poe, 2010). Further, variation in claws may not reflect ecological differences on such a minor scale. In order to stay attached to a vertical perch, the forces produced by pushing the left and right limbs in to the perch must be 90 degrees or less in order to support body weight. Claws are known to help reduce this effective angle (Biewener, 2003; Cartmill, 1974). However, at perch diameters less than a body width, where most of our species were observed, claws are not necessary as the angle between limbs is much less than 90. Thus, a non-existent relationship between perch diameter and claw morphologies agrees with theory.

Claw characters may prove important in ecological contexts not reported in this study. In particular, claw height improves attachment ability on rough surfaces (Zani, 2000). Thus, we predict that improved claw morphology (height, length, curvature, and sharpness) is also correlated with surface roughness. This prediction is opposite that which is known in adhesive toepads, which function best on smooth surfaces (Vanhooydonck et al., 2005). Further, the functional interaction of claw and substrate is unknown. Claws may be used to actively pierce substrates, or attach solely due to passive friction. If claws are used to pierce, our results may be

1
2
3
4 304 confounded by the wearing-down of the claw structure, as observed in artificial
5
6 305 claw systems (Provancher et al., 2004). Use of the claw is likely to vary with size and
7
8 306 morphology of the vertebrate, as well as locomotor style. For example, birds show a
9
10 307 variety of toe organizations – varying from one to two opposing toes in various
11
12 308 configurations. Alongside the differences in muscular arrangement of the flexors
13
14 309 and extensors, this variation may influence whether the claw is actively piercing or
15
16 310 passively attaching to a substrate. Body size and shape has the potential to further
17
18 311 confound claw use. For example, in four-legged vertebrates, the smaller species
19
20 312 tend to have a more sprawled posture, while larger species are more upright
21
22 313 (Biewener, 1989). The distance and angle between opposing limbs likely affects the
23
24 314 effective angle of the claw relative to the substrate – especially on substrates with a
25
26 315 limited diameter (Cartmill, 1974). Behavioral and ecological data in lizards would
27
28 316 shed light on claw use, body size, and locomotion, yet are lacking for most of the
29
30 317 species studied here. For studies outside of *Anolis*, understanding body size, posture,
31
32 318 and foot morphology will be crucial to compare claw morphologies across taxa.

33
34 319 Lastly, although not the focus of our study, we found that mainland anoles
35
36 320 studied here show similar ecological patterns to island species. In island species,
37
38 321 toepad force production positively correlates with perch height (Elstrott and
39
40 322 Irschick, 2004). This potential trend was unknown in mainland species, particularly
41
42 323 because mainland species have dramatically smaller adhesive toepads (Macrini et al.,
43
44 324 2003). Whether or not this difference in size indicates mainland species have less
45
46 325 clinging ability remains to be tested, as island studies are not comparable to our
47
48 326 data and appear more variable in sampling technique. Moreover, our study focused
49
50 327 on a single digit, while previous studies measured force production in the 10
51
52 328 forelimb digits (Irschick et al., 1996). Additionally, mainland species might
53
54 329 experience more diversity in textures and selection for claw characters might not be
55
56 330 as strong as in an island setting. Understanding potential differences and similarities
57
58 331 between clades is of particular interest, as island and mainland species appear to
59
60 332 have different patterns of morphological variation (Schaad and Poe, 2010; Velasco
61
62 333 and Herrel, 2007).

Further studies exploring both mainland and island species in tandem are necessary to elucidate how toepad morphologies differ between populations. Such studies must use caution when collecting or comparing adhesion performance data to previously published values. Our data were collected in the field and may have been influenced by environmental conditions including humidity, which may reduce (Niewiarowski et al., 2008; Stark et al., 2012) or increase (Chen and Gao, 2010; Pesika et al., 2009; Prowse et al., 2011; Puthoff et al., 2010) clinging ability in biological adhesives. In particular, high humidity environments change the mechanical properties of the microscopic setae, increasing adhesion abilities (Prowse et al., 2011; Puthoff et al., 2010). Similarly, increased stress of being sampled in the field may impair performance. Further studies across *Anolis* with standardized methods, alongside increased sample sizes, will improve our understanding of relationships among pad, claw, and habitat use.

Here we have shown the ecological importance of claw morphology. Anoles found higher in the forest have longer and higher claws, with trends toward decreased claw curvature and increased claw tip angle. These results indicate the importance of claws to habitat use. As almost all vertebrates have claws, this character is ripe for further study. Our study also indicates that claws and adhesive toepads have coevolved in *Anolis*. However, they likely serve partially differing functions; a hypothesis that requires further research on the interaction between the toe and natural surfaces. Understanding this interaction will entail an integration of micro-scale studies of single foot-hairs (Autumn, 2006; Autumn et al., 2000; Liang et al., 2000) and claw tips (Dai et al. 2002) with macro-scale studies of the interaction between a variety of natural and artificial surfaces with whole claws (Provancher et al. 2004) and toepads.

Acknowledgements

Field work was arranged through the Organization for Tropical Studies, Duke University, Guido Saborío from Amigos de Osa, and the Smithsonian Tropical Research Institute, under research permit numbers SINAC 123-2008 del Ministerio del Ambiente y Energia – Sistema Nacional de areas de Conservacion and SE/A-64-09 del Autoridad Nacional del Ambiente de Panama. The authors thank Shane Campbell-Staton, Kristi Fenstermacher, Hannah Frank, Martha Muñoz, Stephane Montuelle, Paul VanMiddlesworth, and Katharina Wollenberg for help in the field. For access to specimens, we thank Jose Rosado, Luke Mahler (MCZ, Harvard University), Steve Poe, and Tom Giermakowski (MSB, University of New Mexico). Greta Binford provided feedback on a previous draft. We thank four anonymous reviewers. Use of animals in this study was approved by Lewis & Clark College IACUC (ACUP-2008-001). This research was funded by Sigma Xi (G200803150489 to KEC), Lewis & Clark College SAAB (to KEC), and the National Science Foundation (NSF-NIRT 0304730, NSF-IOS-0847953, NSF_NBM-0900723 and Special Creativity Award to KA; NSF DEB-0444763 to JBL).

Figures

Figure 1. *Anolis sagrei* hanging on a glass slide by a single toe.

Figure 2. Morphological characteristics determined for each digit. Claw height = length of line A; Claw Length = total length of segments C+D; Claw tip angle = Θ , Claw curvature = $57.296 * (2 * \arcsine(((2 * C^2 * D^2) + (2 * B^2 * D^2) + (2 * B^2 * C^2) \pm B^4 \pm C^4 \pm D^4)^{0.5} / (2 * C * D)))$ (modified from Zani, 2000)

Figure 3. Phylogenetic relationships of the 12 species in this study, modified from Nicholson et al. (2005). Branch lengths do not reflect distance.

Figure 4. Independent contrasts showing how toepad force production is correlated with claw height (A) and claw length (B), suggesting that both features have co-evolved in mainland *Anolis* lizards.

Figure 5. Independent contrasts showing that toepad force production is correlated with perch height in mainland *Anolis* lizards, illustrating that both traits have coevolved.

Figure 6. Claw morphological characters for species classified as arboreal (average perch height > 1 meter) and non-arboreal (average perch height < 1 meter). Claw curvature (a) and tip angle (b) are not statistically significantly different, but claw height (c) and length (d) are significantly different between groups.

Table 1. Morphological and performance characters measured for each species. All measurements were taken from the fourth (longest) digit of the right hind foot. Morphological measurements reflect museum specimens (supplement 1), and performance measures were taken from wild-caught individuals.

References

- Arnold, S.J., 1983. Morphology, performance, and fitness. *American Zoologist* 23, 347-361.
- Autumn, K., 2006. Properties, principles, and parameters of the gecko adhesive system, in: Smith, A., Callow, J. (Eds.), *Biological Adhesives*. Springer Verlag, Berlin Heidelberg.
- Autumn, K., Liang, Y.A., Hsieh, S.T., Zesch, W., Chan, W.P., Kenny, T.W., Fearing, R., Full, R.J., 2000. Adhesive force of a single gecko foot-hair. *Nature* 405, 681-685.
- Autumn, K., Sitti, M., Liang, Y.C.A., Peattie, A.M., Hansen, W.R., Sponberg, S., Kenny, T.W., Fearing, R., Israelachvili, J.N., Full, R.J., 2002. Evidence for van der Waals adhesion in gecko setae. *Proceedings of the National Academy of Sciences of the United States of America* 99, 12252-12256.
- Biewener, A.A., 1989. Scaling body support in mammals - limb posture and muscle mechanics. *Science* 245, 45-48.
- Biewener, A.A., 2003. *Animal Locomotion*. Oxford University Press, Oxford, UK.
- Cartmill, M., 1974. Pads and claws in arboreal locomotion, in: Jenkins, F.A. (Ed.), *Primate Locomotion*. Academic Press, New York.
- Castañeda, M., de Queiroz, K., 2013. Phylogeny of the dactyloa clade of *Anolis* lizards: New insights from combining morphological and molecular data. *Bulletin of the Museum of Comparative Zoology* 160, 345-398.
- Chen, B., Gao, H., 2010. An alternative explanation of the effect of humidity in gecko adhesion: Stiffness reduction enhances adhesion on a rough surface. *International Journal of Applied Mechanics* 2, 1-9.
- Dai, Z.D., Gorb, S.N., Schwarz, U., 2002. Roughness-dependent friction force of the tarsal claw system in the beetle *Pachnoda marginata* (Coleoptera, Scarabaeidae). *Journal of Experimental Biology* 205, 2479-2488.
- Elstrott, J., Irschick, D.J., 2004. Evolutionary correlations among morphology, habitat use and clinging performance in Caribbean *Anolis* lizards. *Biological Journal of the Linnean Society* 83, 389-398.
- Feduccia, A., 1993. Evidence from claw geometry indicating arboreal habits of *Archaeopteryx*. *Science* 259, 790-793.
- Felsenstein, J., 1985. Phylogenies and the comparative method. *American Naturalist* 125, 1-15.
- Grafen, A., 1989. The phylogenetic regression. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* 326, 119-157.
- Harmon, L.J., Weir, J.T., Brock, C.D., Glor, R.E., Challenger, W., 2008. GEIGER: investigating evolutionary radiations. *Bioinformatics* 24, 129-131.
- Irschick, D.J., Austin, C.C., Petren, K., Fisher, R.N., Losos, J.B., Ellers, O., 1996. A comparative analysis of clinging ability among pad-bearing lizards. *Biological Journal of the Linnean Society* 59, 21-35.
- Irschick, D.J., Vitt, L.J., Zani, P.A., Losos, J.B., 1997. A comparison of evolutionary radiations in mainland and Caribbean *Anolis* lizards. *Ecology* 78, 2191-2203.

- Liang, Y., Autumn, K., Hsieh, S., Zesch, W., Chan, W., Fearing, R., Full, R., Kenny, T., 2000. Adhesion force measurements on single gecko setae. Technical Digest of the 2000 Solid-State Sensor and Actuator Workshop, 33-38.
- Losos, J.B., 2009. Lizards in an Evolutionary Tree: ecology and adaptive radiation of anoles. University of California Press, Berkeley, CA.
- Macrini, T.E., Irschick, D.J., Losos, J.B., 2003. Ecomorphological differences in toepad characteristics between mainland and island anoles. *Journal of Herpetology* 37, 52-58.
- Maddin, H.C., Reisz, R.R., 2007. Histological microstructure of the claw of the African clawed frog, *Xenopus laevis* (Anura : Pipidae). *Journal of Morphology* 268, 1101-1101.
- Mahendra, B.C., 1941. Contributions to the bionomics, anatomy, reproduction and development of the Indian house gecko *Hemidactylus faviviridis*. Ruppel Part !!. The problem of locomotion. *Proc. Indian Acad. Sci. Sect. B* 13, 288-306.
- Nicholson, K.E., Glor, R.E., Kolbe, J.J., Larson, A., Hedges, S.B., Losos, J.B., 2005. Mainland colonization by island lizards. *Journal of Biogeography* 32, 929-938.
- Niewiarowski, P.H., Lopez, S., Ge, L., Hagan, E., Dhinojwala, A., 2008. Sticky gecko feet: The role of temperature and humidity. *Plos One* 3.
- Persson, B.N.J., Gorb, S., 2003. The effect of surface roughness on the adhesion of elastic plates with application to biological systems. *Journal of Chemical Physics* 119, 11437-11444.
- Pesika, N.S., Zeng, H.B., Kristiansen, K., Zhao, B.X., Tian, Y., Autumn, K., Israelachvili, J., 2009. Gecko adhesion pad: a smart surface? *Journal of Physics-Condensed Matter* 21.
- Provancher, W.R., Clark, J.E., Geisler, B., Cutkosky, M.R., 2004. Towards penetration-based clawed climbing, 7th International Conference on Climbing and Walking Robots (CLAWAR 2004), Madrid, Spain.
- Prowse, M.S., Wilkinson, M., Puthoff, J.B., Mayer, G., Autumn, K., 2011. Effects of humidity on the mechanical properties of gecko setae. *Acta Biomaterialia* 7, 733-738.
- Puthoff, J.B., Prowse, M.S., Wilkinson, M., Autumn, K., 2010. Changes in materials properties explain the effects of humidity on gecko adhesion. *Journal of Experimental Biology* 213, 3699-3704.
- Russell, A., Johnson, M., 2013. Between a rock and a soft place: microtopograph of the locomotor substrate and the morphology of the setal fields of Namibian day geckos (Gekkota: Gekkonidae: *Rhoptropus*). *Acta Zoologica*.
- Schaad, E.W., Poe, S., 2010. Patterns of ecomorphological convergence among mainland and island *Anolis* lizards. *Biological Journal of the Linnean Society* 101, 852-859.
- Stark, A.Y., Sullivan, T.W., Niewiarowski, P.H., 2012. The effect of surface water and wetting on gecko adhesion. *Journal of Experimental Biology* 215, 3080-3086.
- Tulli, M.J., Cruz, F.B., Herrel, A., Vanhooydonck, B., Abdala, V., 2009. The interplay between claw morphology and microhabitat use in neotropical iguanian lizards. *Zoology* 112, 379-392.

- 1
2
3
4 540 Vanhooydonck, B., Andronescu, A., Herrel, A., Irschick, D.J., 2005. Effects of substrate
5 541 structure on speed and acceleration capacity in climbing geckos. Biological
6 542 Journal of the Linnean Society 85, 385-393.
7
8 543 Velasco, J.A., Herrel, A., 2007. Ecomorphology of *Anolis* lizards of the Choco' region in
9 544 Colombia and comparisons with Greater Antillean ecomorphs. Biological Journal
10 545 of the Linnean Society 92, 29-39.
11
12 546 Warheit, K.I., Forman, J.D., Losos, J.B., Miles, D.B., 1999. Morphological diversification
13 547 and adaptive radiation: A comparison of two diverse lizard clades. Evolution 53,
14 548 1226-1234.
15
16 549 Williams, E.E., 1983. Ecomorphs, faunas, island size, and diverse end points in island
17 550 radiations of *Anolis*. , in: Huey, R.B., Pianka, E.R., Schoener, T.W. (Eds.), Lizard
18 551 ecology: Studies of a model organism. Harvard University Press, Cambridge, MA,
19 552 pp. 132-151.
20
21 553 Wollenberg, K.C., Wang, I.J., Glor, R.E., Losos, J.B., 2013. Determinism in the
22 554 diversification of hispaniolan trunk ground anoles (*Anolis cybotes* species
23 555 complex). Evolution.
24
25 556 Zani, P.A., 2000. The comparative evolution of lizard claw and toe morphology and
26 557 clinging performance. Journal of Evolutionary Biology 13, 316-325.
27 558
28 559
29 560
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Species	n (morphology)	n (performance)	SVL (mm)	Mass (g)	Claw height (mm)	Claw Length (mm)	Claw Curvature (degrees)	Claw Tip Angle (degrees)	Toepad Area (mm2)	Lamellae Number	Toepad Force (N)
<i>Anolis auratus</i>	11	6	42.66	1.35	0.32 ± 0.01	0.52 ± 0.03	40.13 ± 1.57	36.34 ± 2.99	1.06 ± 0.07	10 ± 0.23	0.0084 ± 0.00081
<i>Anolis biporcatus</i>	13	3	83.18	12.88	0.79 ± 0.02	1.10 ± 0.04	32.5 ± 0.39	34.39 ± 0.91	7.23 ± 0.34	18.7 ± .3	0.012 ± 0.022
<i>Anolis capito</i>	6	2	75.86	11.22	0.70 ± 0.05	0.90 ± 0.25	35.41 ± 0.91	29.06 ± 2.14	3.60 ± 0.39	12.3 ± 0.33	0.051 ± 0.021
<i>Anolis cupreus</i>	5	4	42.66	1.62	0.32 ± 0.02	0.35 ± 0.10	36.14 ± 1.33	31.60 ± 1.37	1.20 ± 0.23	9.0 ± 0.41	0.019 ± 0.0016
<i>Anolis frenatus</i>	7	9	93.33	17.38	1.14 ± 0.04	1.69 ± 0.1	33.07 ± 0.86	34.28 ± 0.98	18.60 ± 1.90	23.38 ± 1.07	0.15 ± 0.012
<i>Anolis humilis</i>	5	5	31.62	0.91	0.34 ± 0.05	0.52 ± 0.08	34.67 ± 1.03	28.57 ± 1.13	1.32 ± 0.26	10.8 ± 1.11	0.011 ± 0.0011
<i>Anolis lemurinus</i>	5	1	51.29	2.95	0.40 ± 0.01	0.50 ± 0.13	34.24 ± 0.80	35.52 ± 1.07	3.04 ± 0.27	14.0 ± 0.41	0.034 ± 0.0013
<i>Anolis limifrons</i>	7	5	38.02	0.95	0.31 ± 0.02	0.42 ± 0.03	34.07 ± 0.85	29.13 ± 1.64	1.53 ± 0.15	11.0 ± 0.49	0.0073 ± 0.0015
<i>Anolis lionotus</i>	10	7	64.57	5.50	0.46 ± 0.05	0.73 ± 0.11	35.47 ± 0.54	29.50 ± 1.35	3.44 ± 0.52	14 ± 0.24	0.020 ± 0.0013
<i>Anolis pentaprion</i>	9	2	48.98	2.45	0.43 ± 0.03	0.51 ± 0.03	32.91 ± 0.94	36.76 ± 1.27	2.83 ± 0.24	16.9 ± 0.28	0.092 ± 0.012
<i>Anolis poecilopus</i>	6	9	61.66	5.01	0.51 ± 0.05	0.83 ± 0.09	35.67 ± 0.09	29.27 ± 0.66	4.20 ± 0.90	14.4 ± 0.68	0.032 ± 0.0014
<i>Anolis polylepis</i>	6	4	44.67	1.74	0.38 ± 0.02	0.34 ± 0.14	33.31 ± 1.59	31.89 ± 1.19	1.81 ± 0.11	11.8 ± 0.20	0.017 ± 0.0016

Table 1

Species	n (morphology)	n (performance)	SVL (mm)	Mass (g)	Claw height (mm)	Claw Length (mm)
<i>Anolis auratus</i>	11	6	42.66	1.35	0.32 ± 0.01	0.52 ± 0.03
<i>Anolis biporcatus</i>	13	3	83.18	12.88	0.79 ± 0.02	1.1 ± 0.04
<i>Anolis capito</i>	6	2	75.86	11.22	0.70 ± 0.05	0.90 ± 0.25
<i>Anolis cupreus</i>	5	4	42.66	1.62	0.32 ± 0.02	0.35 ± 0.10
<i>Anolis frenatus</i>	7	9	93.33	17.38	1.14 ± 0.04	1.69 ± 0.1
<i>Anolis humilis</i>	5	5	31.62	0.91	0.34 ± 0.05	0.52 ± 0.08
<i>Anolis lemurinus</i>	5	1	51.29	2.95	0.40 ± 0.01	0.50 ± 0.13
<i>Anolis limifrons</i>	7	5	38.02	0.95	0.31 ± 0.02	0.42 ± 0.03
<i>Anolis lionotus</i>	10	7	64.57	5.50	0.46 ± 0.05	0.73 ± 0.11
<i>Anolis pentaprion</i>	9	2	48.98	2.45	0.43 ± 0.03	0.51 ± 0.03
<i>Anolis poecilopus</i>	6	9	61.66	5.01	0.51 ± 0.05	0.83 ± 0.09
<i>Anolis polylepis</i>	6	4	44.67	1.74	0.38 ± 0.02	0.34 ± 0.14

Figure01
[Click here to download high resolution image](#)



Figure02





Figure03







Figure05





Figure06



e-component

[Click here to download e-component: appendix.xlsx](#)